

Coset conformal field theories with abelian isotropy groups.

A.V.Bratchikov*

Kuban State Technological University,
2 Moskovskaya Street, Krasnodar, 350072, Russia.

November, 1998

Abstract

Conformal field theories based on $g/u(1)^d$ coset constructions where g is a reductive algebra are studied. It is shown that the theories are equivalent to constrained WZNW models for g . Generators of extended symmetry algebras and primary fields are constructed.

In a recent paper [1] currents, primary fields and generators of discrete symmetry groups of $g/u(1)^d$, $d = 1, \dots, \text{rank } g$, coset conformal field theories [2, 3] for simple Lie algebras g were constructed. The construction is based on operator quantization of the WZNW theory for g with the constrained affine Lie algebra $\hat{u}(1)^d$ associated with $u(1)^d$. In this article we consider the $g/u(1)^d$ coset theory where g is a direct sum of simple and abelian Lie algebras. Using a version of the generalized canonical quantization method [4, 1] we show that this theory is equivalent to a constrained WZNW theory and find parafermionic currents and primary fields. They are expressed in terms of the initial fields and free bosons. The theory has a discrete symmetry group which is generated by an abelian subalgebra of g .

Studies of such theories were initiated in [5] where $su(2)/u(1)$ parafermionic algebra was constructed and it was shown that the corresponding theory describes a critical system.

*bratchikov@kubstu.ru

Let us consider the WZNW theory [6, 7] for $g = g_1 + g_2 + \dots + g_M$, where g_μ are simple Lie algebras for $1 \leq \mu \leq M'$ and $g_\mu = u(1)^{r_\mu}$ for $M' + 1 \leq \mu \leq M$. We treat only the holomorphic part of the theory. Let k_μ be the central charge of the affine Lie algebra $\hat{g}_\mu$ associated with g_μ .

The affine Lie algebra $\hat{g} = \hat{g}_1 + \hat{g}_2 + \dots + \hat{g}_M$ is generated by elements of the form

$$E(z) = \sum_{\mu'=1}^{M'} \sum_{\alpha \in \Delta_{\mu'}} b_{\mu'}^\alpha E_{\mu'}^\alpha(z) + \sum_{\mu''=M'+1}^M \sum_{s=1}^{r_{\mu''}} b_{\mu''}^s H_{\mu''}^s(z) \quad (1)$$

where $\Delta_{\mu'}$ is the set of the roots and $E_{\mu'}^\alpha(z)$ are step generators of $g_{\mu'}$, $H_{\mu''}^s(z)$ are generators of $\hat{u}(1)^{r_{\mu''}}$, $b = (b_{\mu'}^\alpha, b_{\mu''}^s)$ are numbers.

Let $g_{\mu'}$ be simply-laced. In this case the currents $E_{\mu'}^\alpha(z)$, $H_{\mu''}^s(z)$ and the energy-momentum tensors $L_\mu(z)$ associated with $\hat{g}_\mu$ via Sugawara construction can be expressed in terms of the bosonic fields

$$\varphi_\mu^{sj}(z) = x_\mu^{sj} - i a_{\mu 0}^{sj} \log z + i \sum_{n \neq 0} \frac{a_{\mu n}^{sj}}{n} z^{-n}, \quad (2)$$

where $s = 1, \dots, r_\mu$, $r_{\mu'} \equiv \text{rank } g_{\mu'}$, $j = 1, \dots, k_\mu$, $\mu = 1, \dots, M$ and nonvanishing commutators of the modes are given by

$$[x_\mu^{sj}, a_{\mu 0}^{sj}] = i, \quad [a_{\mu n}^{sj}, a_{\mu(-n)}^{sj}] = n. \quad (3)$$

The bosonic construction for $E_{\mu'}^\alpha(z)$ and $H_{\mu''}^s(z)$ reads [8]

$$E_{\mu'}^\alpha(z) = \sum_{j=1}^{k_{\mu'}} : e^{i\alpha \cdot \varphi_{\mu'}^j(z)} : c_{\mu' \alpha}^j, \quad H_{\mu''}^s(z) = \sum_{j=1}^{k_{\mu''}} i \partial_z \varphi_{\mu''}^{sj}(z). \quad (4)$$

where $::$ denotes normal ordering with respect to the modes of the bosons, $\alpha^2 = 2$ and $c_{\mu' \alpha}^j$ is a cocycle operator. The bosonic construction for $L_{\mu'}$ was obtained in [10]. Let $H_{\mu'}^s(z)$, $1 \leq s \leq r_{\mu'}$ be the currents of $\hat{g}_{\mu'}$ which are defined by

$$H_{\mu'}^s(z) = \sum_{j=1}^{k_{\mu'}} i \partial_z \varphi_{\mu'}^{sj}(z). \quad (5)$$

Let $G^\lambda(z)$, $\lambda = (\lambda_\mu^s)$, $s = 1, \dots, r_\mu$, $\mu = 1, \dots, M$, be the primary field of $\hat{g}$ and the Virasoro algebra generated by $L^g = \sum_{\mu=1}^M L_\mu$ which satisfies the

equations

$$\begin{aligned}
H_{\mu n}^s G^\lambda &= L_{\mu n} G^\lambda = 0, & \text{for } n > 0, \\
H_{\mu 0}^s G^\lambda &= \lambda_\mu^s G^\lambda, \\
L_o^g G^\lambda &= \Delta(\lambda) G^\lambda, & L_{-1}^g G^\lambda = \partial_z G^\lambda,
\end{aligned} \tag{6}$$

where $H_{\mu n}^s$ and $L_{\mu n}$ are modes of the corresponding fields: $H_\mu^s(z) = \sum_n H_{\mu n}^s z^{-n-1}$
 $L_\mu(z) = \sum_n L_{\mu n} z^{-n-2}$.

Let us consider the $\hat{u}^d(1)_k$, $k = \sum_{\mu=1}^M k_\mu$, subalgebra of $\hat{g}$ generated by the currents

$$H^A(z) = \sum_{\mu=1}^M H_\mu^A(z) \tag{7}$$

where $A = 1, \dots, d$, $d \leq r_\mu$, $\mu = 1, \dots, M$. We shall consider the WZNW theory subject to the constraints

$$H^A(z) \approx 0, \tag{8}$$

This equation can be rewritten in the form

$$H_n^A \approx 0, \tag{9}$$

where $H_n^A = \sum_{\mu=1}^M H_{\mu n}^A$.

To quantize the system we put into correspondence with the constraints the bosonic fields

$$\phi^A(z) = q^A - i\tilde{a}_0^A \log z + i \sum_{n \neq 0} \frac{\tilde{a}_n^{Aj}}{n} z^{-n}, \tag{10}$$

where nonvanishing commutators of the modes are given by

$$[q^A, \tilde{a}_0^A] = ik, \quad [\tilde{a}_n^A, \tilde{a}_{-n}^A] = nk. \tag{11}$$

According to a version of the generalized canonical quantization method [4, 1] we replace the constraints (8) by the effective abelian constraints

$$\tilde{H}^A(z) = H^A(z) + \partial \phi^A(z) \approx 0. \tag{12}$$

It is easy to see that

$$\tilde{H}^A(z)\tilde{H}^B(w) = 0. \quad (13)$$

The current $E(z)$ (1) is replaced by

$$\mathcal{E}(z) = \sum_{\mu'=1}^M \sum_{\alpha \in \Delta_{\mu'}} b_{\mu'}^{\alpha} \mathcal{E}_{\mu'}^{\alpha}(z) + \sum_{\mu''=M'+1}^M \sum_{s=1}^{r_{\mu''}} b_{\mu''}^s \mathcal{H}_{\mu''}^s(z), \quad (14)$$

$$\mathcal{E}_{\mu'}^{\alpha}(z) = E_{\mu'}^{\alpha}(z) : e^{\frac{1}{k}\tilde{\alpha} \cdot \phi(z)} :, \quad (15)$$

$$\mathcal{H}_{\mu''}^s(z) = H_{\mu''}^s(z) + \frac{k_{\mu''}}{k} \sum_{A=1}^d \delta^{sA} \partial \phi^A(z), \quad (16)$$

where $\tilde{\alpha} = (\alpha^A)$. The new current commute with $\tilde{H}^A(z)$

$$\tilde{H}^A(z)\mathcal{E}(w) = 0. \quad (17)$$

The $g/u(1)^d$ energy-momentum tensor $K = L^g - L^{u(1)^d}$, where

$$L^{u(1)^d} = \frac{1}{2k} \sum_{A=1}^d : (H^A)^2 :, \quad (18)$$

also commutes with $\tilde{H}^A(z)$ [3]

$$\tilde{H}^A(z)K(w) = 0. \quad (19)$$

The operator product expansion of K with $\mathcal{E}$ is given by

$$\begin{aligned} K(z)\mathcal{E}(w) &= \frac{1}{(z-w)^2} \left(\mathcal{E}(w) - \frac{1}{2k} \sum_{\mu'=1}^{M'} \sum_{\alpha \in \Delta_{\mu'}} \tilde{\alpha}^2 b_{\mu'}^{\alpha} \mathcal{E}_{\mu'}^{\alpha}(w) \right) \\ &+ \frac{1}{z-w} \partial_w \mathcal{E}(w) - \frac{1}{k} \delta \mathcal{E}(b, \tilde{H}^A), \end{aligned} \quad (20)$$

where

$$\begin{aligned} \delta \mathcal{E}(b, \tilde{H}^A) &= \frac{1}{(z-w)^2} \sum_{\mu''=M'+1}^M \sum_{A=1}^d k_{\mu''} b_{\mu''}^A \tilde{H}^A(w) \\ &+ \frac{1}{(z-w)} \left(\sum_{\mu'=1}^{M'} \sum_{\alpha \in \Delta_{\mu'}} b_{\mu'}^{\alpha} \tilde{\alpha} \cdot \tilde{H}(w) \mathcal{E}^{\alpha}(w) \right. \\ &\left. + \sum_{\mu''=M'+1}^M \sum_{A=1}^d k_{\mu''} b_{\mu''}^A \partial_w \tilde{H}^A(w) \right). \end{aligned}$$

Note that $\delta\mathcal{E}(b, 0) = 0$.

If $b_{\mu'}^\alpha$ satisfy the equations

$$\tilde{\alpha}^2 b_{\mu'}^\alpha = \nu^2 b_{\mu'}^\alpha, \quad (21)$$

where ν^2 is a constant, then eq.(20) can be rewritten in the form

$$K(z)\mathcal{E}(w) = \frac{\Delta\mathcal{E}(w)}{(z-w)^2} + \frac{\partial_w\mathcal{E}(w)}{z-w} - \delta\mathcal{E}(b, \tilde{H}^A), \quad (22)$$

where

$$\Delta = 1 - \frac{1}{2k}\nu^2.$$

Let Ω be the algebra generated by $\mathcal{E}(z)$ and $K(z)$. Let us define the following set of the fields

$$\Upsilon = (U \in \Omega | U|_{\tilde{H}^A=0} = 0). \quad (23)$$

Using Wick theorem one can check that for arbitrary $U \in \Upsilon$ and $X \in \Omega$

$$UX \in \Upsilon, \quad XU \in \Upsilon. \quad (24)$$

Hence Υ is an ideal of Ω and the quotient Ω/Υ is an algebra. We shall denote by $\{X(z)\}$ the coset represented by the field $X(z)$.

The coset field $\{K(z)\}$ satisfies the $g/u(1)^d$ Virasoro algebra:

$$\{K(z)\}\{K(w)\} = \frac{c_{g/u(1)^d}}{2(z-w)^4} + \frac{2\{K(w)\}}{(z-w)^2} + \frac{\partial_w\{K(w)\}}{z-w}, \quad (25)$$

where $c_{g/u(1)^d} = c_g - d$.

It follows from eq. (22) that $\{\mathcal{E}(z)\}$ is a primary field of the $g/u(1)^d$ Virasoro algebra:

$$\{K(z)\}\{\mathcal{E}(w)\} = \frac{\Delta\{\mathcal{E}(w)\}}{(z-w)^2} + \frac{\partial_w\{\mathcal{E}(w)\}}{z-w}. \quad (26)$$

The theory has the discrete symmetry group which is generated by $H_{\mu 0}^s$, $s = 1, \dots, r_\mu$, $\mu = 1, \dots, M$.

The initial primary field G^λ is replaced by

$$\mathcal{G}^\lambda(z) = G^\lambda(z) : e^{\frac{1}{k}\tilde{\lambda} \cdot \phi(z)} :, \quad (27)$$

where $\tilde{\lambda}^A = \sum_{\mu=1}^M \lambda_\mu^A$. This field satisfies the following equations

$$\begin{aligned} \tilde{H}_n^A \mathcal{G}^\lambda &= 0, \quad \text{for } n \geq 0, & K_n \mathcal{G}^\lambda &= 0, \quad \text{for } n > 0, \\ K_o \mathcal{G}^\lambda &= \Delta(\lambda, \tilde{\lambda}) \mathcal{G}^\lambda, & K_{-1} \mathcal{G}^\lambda &= \partial_z \mathcal{G}^\lambda - \frac{1}{k} \tilde{\lambda} \cdot \tilde{H}_{-1} \mathcal{G}^\lambda, \end{aligned} \quad (28)$$

where

$$\Delta(\lambda, \tilde{\lambda}) = \Delta(\lambda) - \frac{1}{2k} \tilde{\lambda}^2. \quad (29)$$

Let T be the space which is obtained by applying the currents of Ω repeatedly to the field $\mathcal{G}^\lambda$. Let

$$V = (\mathcal{G} \in T | \mathcal{G}|_{\tilde{H}^A=0} = 0). \quad (30)$$

The space V is an invariant subspace with respect to the algebra Ω . It follows from eq.(28) that $\{\mathcal{G}^\lambda\} \in T/V$ which is represented by $\mathcal{G}^\lambda$ is a primary field of the coset Virasoro algebra.

The space T/V can be decomposed according to the transformation properties under the algebra generated by $H_{\mu 0}^s, \mu = 1, \dots, M, s = 1, \dots, r_\mu$. The field $\{\mathcal{G}^\lambda\}$ belongs to the subspace with the charges λ_μ^s :

$$H_{\mu 0}^s \{\mathcal{G}^\lambda\} = \lambda_\mu^s \{\mathcal{G}^\lambda\}. \quad (31)$$

The theory can be formulated in terms of the initial operators x_μ^{sj} and $a_{\mu n}^{sj}$. To show this we first note that the coset $\{\mathcal{E}(z)\}$ can be represented by the field

$$\tilde{\mathcal{E}}(z) = \mathcal{E}(z)|_{\tilde{a}_n^A = i a_n^A}. \quad (32)$$

It follows from (26) that $\tilde{\mathcal{E}}$ satisfies the equation

$$K(z) \tilde{\mathcal{E}}(w) = \frac{\Delta_\alpha \tilde{\mathcal{E}}(w)}{(z-w)^2} + \frac{\partial_w \tilde{\mathcal{E}}(w)}{z-w}. \quad (33)$$

$\tilde{\mathcal{E}}$ still depends on the auxiliary operators q^A . However these operators commute with $x_\mu^{sj}, a_{\mu n}^{sj}$ and hence one can set $q^A = 0$. The field $\mathcal{E}_0 = \tilde{\mathcal{E}}|_{q^A=0}$ satisfies eq.(33) as well as $\tilde{\mathcal{E}}$.

These results can be generalized to non-simply-laced algebras using the vertex operator representation of the associated affine KM algebras [11, 12].

References

- [1] A.V.Bratchikov, $g/u(1)^d$ parafermions from constrained WZNW theories. hep-th/9712143.
- [2] K.Bardakci and M.B.Halpern, *Phys.Rev.* **D3**(1971)2493.
M.B.Halpern, *Phys.Rev.* **D4**(1971)2398.
- [3] P.Goddard, A.Kent and D.Olive, *Phys.Lett.* **152B** (1985) 88.
- [4] I.A.Batalin, E.S.Fradkin and T.Fradkina, *Nucl.Phys.* **B332** (1990) 723.
- [5] V.A.Fateev and A.B.Zamolodchikov, *Sov.Phys.JETP* **82** (1985) 215.
- [6] E.Witten, *Comm.Math.Phys.* **92** (1984) 455.
- [7] V.Knizhnik and A.B.Zamolodchikov, *Nucl.Phys.* **B247** (1984) 83.
- [8] I.B.Frenkel and V.G.Kac, *Inv.Math.* **62**(1980)23.
- [9] G.Segal, *Commun.Math.Phys.* **80** (1981) 301.
- [10] G.V.Dunne, I.G.Halliday and P.Suranyi, *Nucl.Phys.* **B325** (1989) 526.
- [11] P.Goddard, W.Nahm, D.Olive and A.Schwimmer, *Commun.Math.Phys.* **107** (1986) 179.
- [12] D.Bernard and J.Thierry-Mieg, *Commun.Math.Phys.* **111** (1987) 181.